

PROPERTIES OF CERTAIN BILATERAL MOCK THETA FUNCTIONS-V

Mohammad Ahmad

Abstract

AMS Subject Classification:

Primary 33D15, Secondary
11B6

Keywords:

Mock theta functions; bilateral
mock theta functions^{*}; Lerch
transcendent; F -function.

Bilateral mock theta functions were obtained and studied in [9]. We express them in terms of Lerch's transcendental function $f(x, \xi; q, p)$. We also express some bilateral mock theta functions as sum of other mock theta functions. We generalize these functions and show that these generalizations are F_q functions. We give an integral representation for these generalized functions.

Copyright © 2017 International Journals of Multidisciplinary
Research Academy. All rights reserved.

Author correspondence:

Department of Mathematics
National Defence Academy Khadakwasla Pune-411023

1. Introduction: The mock theta functions were first introduced by Ramanujan [3] in his last letter to G. H. Hardy in January 1920. He provided a list of seventeen mock theta functions and labelled them as of third, fifth and seventh order without mentioning the reason for his labelling. Watson [21] added to this set three more third order mock theta functions.

His general definition of a mock theta function is a function $f(q)$ defined by q -series convergent when $|q| < 1$ which satisfies the following two conditions.

(a) For every root ξ of unity, there exists a theta function[†] $\theta_\xi(q)$ such that the difference between $f(q)$ and $\theta_\xi(q)$ is bounded as $q \rightarrow \xi$ radially.

(b) There is no single theta function which works for all ξ i.e. for every theta function $\theta_\xi(q)$ there is some root of unity ξ for which $f(q)$ minus the theta function $\theta_\xi(q)$ is unbounded as $q \rightarrow \xi$ radially.

^{*} In bilateral form summation is taken from $-\infty$ to ∞ .

[†] When Ramanujan refers to theta functions, he means sums, products, and quotients of series of the form $\sum_{n \in \mathbb{Z}} \epsilon^n q^{an^2+bn}$ with $a, b \in \mathbb{Q}$ and $\epsilon = -1, 1$.

Andrews and Hickerson [18] announced the existence of eleven more identities given in the 'Lost' note book of Ramanujan involving seven new functions which they labelled as mock theta functions of order six.

Y. S. Choi [1] has discovered four functions which he called the mock theta function of order ten. B. Gordon and R. J. McIntosh [33] have announced the existence of eight mock theta functions of order eight and R. J. McIntosh [5] has announced the existence of three mock theta functions of order two.

Hikami [16], [17] has introduced a mock theta function of order two, another of order four and two of order eight. Very recently Andrews [19] while studying q -orthogonal polynomials found four new mock theta functions and Bringmann et al [15] have also found two more new mock theta functions but they did not mention the order of their mock theta functions.

Watson [22] has defined four bilateral series, which he has called the 'Complete' or Bilateral forms for four of the ten mock theta functions of order five. Further he has expressed them in terms of the transcendental function $f(x, \xi; q, p)$ studied by M. Lerch [7]. S. D. Prasad [2] in 1970 has defined the 'Complete' or 'Bilateral' forms of the five generalized third order mock theta functions. The 'Complete' sixth order mock theta functions were studied by A. Gupta [34]. Bhaskar Srivastava [29], [30], [31], [32] have studied bilateral mock theta functions of order five, eight, two and new mock theta functions by Andrews [6] and Bringmann et al [15].

Truesdell [28] calls the functions which satisfy the equation $\frac{\partial}{\partial z} F(z, \alpha) = F(z, \alpha + 1)$ as F - functions. He has tried to unify the study of these F -functions. The function which satisfy the q - analogue of the equation $D_{q,z} F(z, \alpha) = F(z, \alpha + 1)$ where $zD_{q,z} F(z, \alpha) = F(z, \alpha) - F(zq, \alpha)$ are called F_q - functions.

Mohammad Ahmad and Shahab Faruq have obtained and studied the following bilateral mock theta functions in [9].

$$f_{0,c7}(q) = \sum_{-\infty}^{\infty} (-1)^n \frac{q^{\left(\frac{7n^2}{2} - \frac{5n}{2}\right)}}{(-q; q)_n}, \quad (1.1)$$

$$f_{1,c7}(q) = \sum_{-\infty}^{\infty} (-1)^n \frac{q^{\left(\frac{7n^2}{2} - \frac{3n}{2}\right)}}{(-q; q)_n}, \quad (1.2)$$

$$F_{0,c7}(q^2) = \sum_{-\infty}^{\infty} (-1)^n \frac{q^{(7n^2-5n)}}{(q; q^2)_n}, \quad (1.3)$$

$$F_{1,c7}(q^4) = \sum_{-\infty}^{\infty} (-1)^n \frac{q^{(14n^2-6n)}}{(q^6; q^4)_n}, \quad (1.4)$$

$$\Psi_{0,c7}(q) = \sum_{-\infty}^{\infty} (-1)^n q^{(3n^2+9n)} (-q; q)_n \quad (1.5)$$

$$\Phi_{1,c7}(q^2) = \sum_{-\infty}^{\infty} (-1)^n q^{(6n^2+12n)} (-q; q^2)_n \tag{1.6}$$

$$\Phi_{0,c7}(q^2) = \sum_{-\infty}^{\infty} (-1)^n \frac{q^{7n^2}}{(-q; q^2)_n} \tag{1.7}$$

$$\Psi_{1,c7}(q) = \sum_{-\infty}^{\infty} (-1)^{n+1} \frac{q^{q\left(\frac{7n^2}{2} + \frac{7n}{2}\right)}}{2(-q; q)_n} \tag{1.8}$$

The paper is divided as follows: In section 2 we list few important definitions. In section 3 we develop certain properties of these functions by expressing some of them as sums of other mock theta functions. We also express these functions in terms of the Lerch transcendental function $f(x, \xi; q, p)$. In section 4 we generalize these functions which are then proved to be F_q functions. We further give an integral representation of these functions.

2. Notation and Definitions:

We use the following q -notation. Suppose q and z are complex numbers and n is an integer. If $n \geq 0$ we define

$$(z)_n = (z; q)_n = \prod_{i=0}^{n-1} (1 - q^i z) \text{ if } n \leq 0 \text{ and } (z)_{-n} = (z; q)_{-n} = \frac{(-z)^{-n} q^{n(n+1)/2}}{\left(\frac{z}{q}\right)_n}$$

and more generally $(z_1, z_2, \dots, z_r; q)_n = (z_1)_n (z_2)_n \dots (z_r)_n$.

For $|q^k| < 1$ let us define

$$(z; q^k)_n = (1 - z)(1 - zq^k) \dots (1 - zq^{k(n-1)}) \quad n \geq 1 \quad (z; q^k)_0 = 1$$

and $(z; q^k)_\infty = \lim_{n \rightarrow \infty} (z; q^k)_n = \prod_{i \geq 0} (1 - q^{ki} z)$ and even more generally,

$$(z_1, z_2 \dots z_r; q^k)_\infty = (z_1; q^k)_\infty \dots (z_r; q^k)_\infty.$$

A basic hypergeometric series ${}_{r+1}\Phi_r$ on base q^k is defined as

$${}_{r+1}\Phi_r \left[\begin{matrix} a_1, a_2, \dots, a_p \\ b_1, b_2, \dots, b_q \end{matrix}; q \quad z \right] = \sum_{n=0}^{\infty} \frac{(a_1, a_2, \dots, a_p; q^k)_n z^n}{(q^k; q^k)_n (b_1, b_2, \dots, b_r; q^k)_n}, \quad (|z| < 1) \tag{2.1}$$

and a bilateral basic hypergeometric series ${}_r\Psi_r$ is defined as

$${}_{r+1}\Psi_r \left[\begin{matrix} a_1, a_2, \dots, a_p \\ b_1, b_2, \dots, b_q \end{matrix}; q \quad z \right] = \sum_{n=-\infty}^{\infty} \frac{(a_1, a_2, \dots, a_p; q^k)_n z^n}{(q^k; q^k)_n (b_1, b_2, \dots, b_r; q^k)_n}, \quad \left(\left| \frac{b_1 \dots b_r}{a_1 \dots a_p} \right| < |z| < 1 \right) \tag{2.2}$$

The Lerch transcendental function $f(x, \xi; q, p)$ is defined by:

$$f(x, \xi; q, p) = \sum_{-\infty}^{\infty} \frac{(pq)^{n^2} (x\xi)^{-2n}}{(-p\xi^{-2}; p^2)_n} \tag{2.3} \quad \text{and by}$$

$$f(x, \xi; q, p) = \sum_{-\infty}^{\infty} (-\xi^2 p; p^2)_n q^{n^2} x^{2n}. \tag{2.4}$$

3. Certain Identities and Their Lerch Representation:

The following identities between the bilateral mock theta functions given in Equations 1.1, 1.5, 1.6, 1.7 and the corresponding mock theta functions may be verified by hypergeometric transformations:

$$f_{0,c_7}(q) = f_{0,7}(q) - 2q^6\Psi_{0,7}(q) \tag{3.1}$$

$$\Psi_{0,c_7}(q) = \Psi_{0,7}(q) - \frac{1}{2q^6}f_{0,7}(q) \tag{3.2}$$

$$\Phi_{0,c_7}(q^2) = \Phi_{0,7}(q^2) - q^6\Phi_{1,7}(q^2) \sum_0^\infty (1 + q^{2n+1}) \tag{3.3}$$

Here $f_{0,7}(q), \Psi_{0,7}(q), \Phi_{0,7}(q^2), \Phi_{1,7}(q^2)$ are the corresponding mock theta functions.

The bilateral mock theta functions defined in Section 1 can be expressed in terms of the Lerch transcendent by means of the following lemma.

Lemma 3.1 For $\epsilon = \pm 1$,

$$\sum_{n=-\infty}^\infty (-1)^n \frac{q^{\alpha n^2} q^{\beta n}}{(\epsilon q^\gamma; q^\delta)_n} = f\left(i(-\epsilon)^{\frac{1}{2}} q^{\frac{2\gamma-2\beta-\delta}{4}}, (-\epsilon)^{\frac{1}{2}} q^{\frac{\delta-2\gamma}{4}}; q^{\frac{2\alpha-\delta}{2}}, q^{\frac{\delta}{2}}\right).$$

and

$$\sum_{n=-\infty}^\infty (-1)^n (-q; q^\gamma)_n q^{\alpha n^2} q^{\beta n} = f\left(iq^{\frac{\beta}{2}}, q^{\frac{2-\gamma}{4}}; q^\alpha, q^{\frac{\gamma}{2}}\right).$$

Proof: The proof follows from direct substitution and use of basic hypergeometric transformations.

As an example we note that $f_{0,c_7}(q) = \sum_{n=-\infty}^\infty (-1)^n \frac{q^{\left(\frac{7n^2}{2} - \frac{3n}{2}\right)}}{(-q; q)_n} = f\left(iq, q^{\frac{1}{4}}; q^3, q^{\frac{1}{2}}\right)$ by taking $\alpha = \frac{7}{2}, \beta = \frac{-3}{2}, \epsilon = -1, \gamma = \delta = 1$ and $\Psi_{0,c_7}(q) = \sum_{n=-\infty}^\infty (-1)^n q^{(3n^2+9n)} (-q; q)_n = f\left(iq^{\frac{9}{2}}, q^{\frac{1}{4}}; q^3, q^{\frac{1}{2}}\right)$ by taking $\alpha = 2, \beta = 9, \gamma = 1$ in the above lemma.

In this way all other bilateral mock theta functions defined by Equations 1.1 to 1.8 can be expressed in terms of the Lerch Transcendental function defined by equations (2.3) and (2.4).

4. Generalisation of Bilateral Mock Theta Functions:

We generalize the functions given by Equations 1.1 to 1.8 by introducing two parameters α, z . For $\alpha = 1, z = 0$ these are reduced to the original functions.

$$f_{0,c_7}(z, \alpha; q) = \frac{1}{(z)_\infty} \sum_{n=-\infty}^\infty (-1)^n \frac{(z)_n q^{\frac{7n^2}{2} + n\alpha - \frac{7n}{2}}}{(-q; q)_n} \tag{4.1}$$

$$f_{1,c_7}(z, \alpha; q) = \frac{1}{(z)_\infty} \sum_{n=-\infty}^\infty (-1)^n \frac{(z)_n q^{\frac{7n^2}{2} + n\alpha - \frac{5n}{2}}}{(-q; q)_n} \tag{4.2}$$

$$F_{0,c_7}(z, \alpha; q^2) = \frac{1}{(z)_\infty} \sum_{-\infty}^{\infty} (-1)^n \frac{(z)_n q^{7n^2 + \alpha - 6n}}{(q; q^2)_n} \quad (4.3)$$

$$F_{1,c_7}(z, \alpha; q^4) = \frac{1}{(z)_\infty} \sum_{-\infty}^{\infty} (-1)^n \frac{(z)_n q^{14n^2 + \alpha - 7n}}{(q^6; q^4)_n} \quad (4.4)$$

$$\Psi_{0,c_7}(z, \alpha; q) = \frac{1}{(z)_\infty} \sum_{-\infty}^{\infty} (z)_n (-1)^n q^{3n^2 + \alpha + 8n} (-q; q)_n \quad (4.5)$$

$$\Phi_{1,c_7}(z, \alpha; q^2) = \frac{1}{(z)_\infty} \sum_{-\infty}^{\infty} (z)_n (-1)^n q^{6n^2 + \alpha + 11n} (-q; q^2)_n \quad (4.6)$$

$$\Phi_{0,c_7}(z, \alpha; q^2) = \frac{1}{(z)_\infty} \sum_{-\infty}^{\infty} (-1)^n \frac{(z)_n q^{7n^2 + \alpha - n}}{(-q; q^2)_n} \quad (4.7)$$

$$\Psi_{1,c_7}(z, \alpha; q) = \sum_{-\infty}^{\infty} (-1)^{n+1} \frac{(z)_n q^{\frac{7n^2}{2} + \alpha + \frac{5n}{2}}}{2(-q; q)_n} \quad (4.8)$$

We now show that these generalized functions are F_q functions.

Theorem 4.1

The functions defined by the Equations 4.1 – 4.8 are F_q functions.

Proof: We give the proof only for $f_{0,c_7}(z, \alpha; q)$. The remaining cases are similar. For $f_{0,c_7}(z, \alpha; q)$ note that

$$\begin{aligned} zD_{q,z} f_{0,c_7}(z, \alpha; q) &= f_{0,c_7}(z, \alpha; q) - f_{0,c_7}(zq, \alpha; q) \\ &= \frac{1}{(z)_\infty} \sum_{-\infty}^{\infty} (-1)^n \frac{(z)_n q^{\frac{7n^2}{2} + \alpha - \frac{7n}{2}}}{(-q; q)_n} \\ &\quad - \frac{1}{(zq)_\infty} \sum_{-\infty}^{\infty} (-1)^n \frac{(zq)_n q^{\frac{7n^2}{2} + \alpha - \frac{7n}{2}}}{(-q; q)_n} \\ &= \frac{1}{(z)_\infty} \sum_{-\infty}^{\infty} (-1)^n \frac{(z)_n q^{\frac{7n^2}{2} + \alpha - \frac{7n}{2}}}{(-q; q)_n} (1 - (1 - zq^n)) \\ &= \frac{z}{(z)_\infty} \sum_{-\infty}^{\infty} (-1)^n \frac{(z)_n q^{\frac{7n^2}{2} + (n+1)\alpha - \frac{7n}{2}}}{(-q; q)_n} \\ &= z f_{0,c_7}(z, \alpha + 1; q) \end{aligned}$$

and hence $f_{0,c_7}(z, \alpha; q)$ is a F_q function.

We now give integral representations of these generalized functions. Jackson (on Page 23 of [17]) defined the q -integral on $(0, \infty)$ by

$$\int_0^{\infty} f(t) d_q t = (1-q) \sum_{n=-\infty}^{\infty} f(q^n) q^n.$$

Now let $f(t) = t^{x-1}(tq; q)_{\infty}$ for some fixed x . We have

$$\begin{aligned} \int_0^{\infty} t^{x-1}(tq; q)_{\infty} d_q t &= (1-q) \sum_{n=-\infty}^{\infty} (q^{n+1}; q)_{\infty} q^{nx} \\ &= (1-q) \frac{(q; q)_{\infty}}{(q^x; q)_{\infty}} \end{aligned}$$

and so

$$\frac{1}{(q^x; q)_{\infty}} = \frac{(1-q)^{-1}}{(q; q)_{\infty}} \int_0^{\infty} t^{x-1}(tq; q)_{\infty} d_q t. \quad (4.9)$$

We now use Equation 4.9 to give integral representations of the F_q functions 4.1 to 4.8. We let $a = q^{\alpha}$ for convenience.

$$f_{0,c7}(q^z, \alpha; q) = \frac{(1-q)^{-1}}{(q; q)_{\infty}} \int_0^{\infty} t^{z-1}(tq; q)_{\infty} f_{0,c7}(0, at; q) d_q t \quad (4.10)$$

$$f_{1,c7}(q^z, \alpha; q) = \frac{(1-q)^{-1}}{(q; q)_{\infty}} \int_0^{\infty} t^{z-1}(tq; q)_{\infty} f_{1,c7}(0, at; q) d_q t \quad (4.11)$$

$$F_{0,c7}(q^z, \alpha; q^2) = \frac{(1-q)^{-1}}{(q; q)_{\infty}} \int_0^{\infty} t^{z-1}(tq; q)_{\infty} F_{0,c7}(0, at; q^2) d_q t \quad (4.12)$$

$$F_{1,c7}(q^z, \alpha; q^4) = \frac{(1-q)^{-1}}{(q; q)_{\infty}} \int_0^{\infty} t^{z-1}(tq; q)_{\infty} F_{1,c7}(0, at; q^4) d_q t \quad (4.13)$$

$$\Psi_{0,c7}(q^z, \alpha; q) = \frac{(1-q)^{-1}}{(q; q)_{\infty}} \int_0^{\infty} t^{z-1}(tq; q)_{\infty} \Psi_{0,c7}(0, at; q) d_q t \quad (4.14)$$

$$\Phi_{1,c7}(q^z, \alpha; q^2) = \frac{(1-q)^{-1}}{(q; q)_{\infty}} \int_0^{\infty} t^{z-1}(tq; q)_{\infty} \Phi_{1,c7}(0, at; q^2) d_q t \quad (4.15)$$

$$\Phi_{0,c7}(q^z, \alpha; q^2) = \frac{(1-q)^{-1}}{(q; q)_{\infty}} \int_0^{\infty} t^{z-1}(tq; q)_{\infty} \Phi_{0,c7}(0, at; q^2) d_q t \quad (4.16)$$

$$\Psi_{1,c7}(q^z, \alpha; q) = \frac{(1-q)^{-1}}{(q; q)_{\infty}} \int_0^{\infty} t^{z-1}(tq; q)_{\infty} \Psi_{1,c7}(0, at; q) d_q t \quad (4.17)$$

Theorem 4.2 Equations 4.10 to 4.17 hold.

Proof: We prove only 4.10. The remaining cases are similar. Writing q^z for z and a for q^{α} in 4.1 we have,

$$\begin{aligned}
f_{0,c7}(q^z, \alpha; q) &= \frac{1}{(q^z; q)_\infty} \sum_{-\infty}^{\infty} (-1)^n \frac{a^n (q^z; q)_n q^{\frac{7n^2}{2} - \frac{7n}{2}}}{(-q; q)_n} \\
&= \sum_{-\infty}^{\infty} (-1)^n \frac{a^n q^{\frac{7n^2}{2} - \frac{7n}{2}}}{(-q; q)_n (q^{n+z}; q)_\infty} \\
&= \frac{(1-q)^{-1}}{(q; q)_\infty} \sum_{-\infty}^{\infty} (-1)^n \frac{a^n q^{\frac{7n^2}{2} - \frac{7n}{2}}}{(-q; q)_n} \int_0^\infty t^{n+z-1} (tq; q)_\infty d_q t \\
&= \frac{(1-q)^{-1}}{(q; q)_\infty} \int_0^\infty t^{z-1} (tq; q)_\infty \sum_{-\infty}^{\infty} (-1)^n \frac{(at)^n q^{\frac{7n^2}{2} - \frac{7n}{2}}}{(-q; q)_n} d_q t \\
&= \frac{(1-q)^{-1}}{(q; q)_\infty} \int_0^\infty t^{z-1} (tq; q)_\infty f_{0,c7}(0, at; q) d_q t
\end{aligned}$$

which completes the proof.

We remark that for $at = q$ the function $f_{0,c7}(0, at; q)$ reduces to the bilateral mock theta function $f_{0,c7}(q)$ defined previously.

Acknowledgement: Support and guidance of Prof O. P. Shukla, Principal NDA Khadakwasla and Prof D P Shukla, Ex Professor of Mathematics, Lucknow University Lucknow is gratefully acknowledged.

References:

- [1] Y. S. Choi, Tenth order mock theta functions in Ramanujan's lost note book, *Inventiones Mathematicae* 136(1999), 497-569.
- [2] S.D. Prasad, Certain extended mock theta functions and generalized basic hypergeometric transformation, *Math Scand* 27(1970), 237-244.
- [3] S. Ramanujan, *Collected Papers*, Cambridge University Press (1927), reprinted by Chelsea New York, 1960.
- [4] R.P. Agarwal, Mock theta functions-An analytical point of view, *Proc. Nat. Acad. Sci. India.* 64(A.1) (1994), 95-106.
- [5] R. J. McIntosh, Second order mock theta functions, *Canadian Mathematical Bulletin* 50 (2)(2007), 284-290.
- [6] N. J. Fine, *Basic hypergeometric series and applications Mathematical Surveys & Mono Graphs*, Providence, R.I Amer. Math. Soc. 27, 1988.
- [7] M. Lerch. , Novanalogierady theta a nektezvltnhypergeometrickradyHeineovy. *Rozpravy* 3(1893), 1–10.
- [8] M Ahmad, *On the Behavior of Bilateral Mock Theta Functions-I*, Algebra and Analysis: Theory and Application, Narosa Publishing House Pvt Ltd, New Delhi, (2015), 259-273.
- [9] M Ahmad and Shahab Faruqi, Some Bilateral Mock Theta Functions and Their Lerch Representations, *The Aligarh Bulletin of Mathematics*, Vol 34, Numbers 1-2(2015) 75-92.

- [10] M Ahmad and Shahab Faruqi, Properties of Certain Bilateral Mock Theta Functions, to appear in Vol 6.(2017) in GAMS Journal of Mathematics and Mathematical Bioscience.
- [11] M Ahmad and Raghieb Nadeem, Properties of Certain Bilateral Mock Theta Functions-II to appear in Vol 10(1),(2018) in Global-Sci.Tech. Journal.
- [12] M Ahmad, "Properties of Certain Bilateral Mock Theta Functions-III" International Journal of Engineering, Science and Mathematics Vol. 6 Issue 6 (2017) 224-232.
- [13] M Ahmad, "Properties of Certain Bilateral Mock Theta Functions-IV" International Journal of Scientific Research in Science and Technology Vol. 3 Issue 7 (2017) 789-793.
- [14] L.A. Dragonette, Some asymptotic formulae for the mock theta series of Ramanujan, Trans. Amer. Math. Soc. **72** (1952), 474-500.
- [15] K. Bringmann, K. Hikami and Jeremy Lovejoy, On the modularity of the unified WRT invariants of certain Seifert manifolds, Advances in Applied Mathematics 46(1) (2011), 86-93.
- [16] K. Hikami, Mock (false) theta functions as quantum invariants, Regular and Chaotic Dynamics, 10 (2005), 509-530.
- [17] K. Hikami, Transformation formulae of the 2nd order mock theta function, Lett. Math. Phys. 75(1) (2006), 93-98.
- [18] G.E. Andrews and D. Hickerson, The sixth order mock theta functions, Adv. Maths 89 (1991), 60-105.
- [19] G. E. Andrews, q-orthogonal polynomials, Roger-Ramanujan identities and mock theta functions, Proc. Steklov Institute of Math. 276(1)(2012), 21-32.
- [20] G. Gasper and M. Rahman, Basic Hypergeometric Series, 2nd ed, Cambridge University Press Cambridge (UK), 2004.
- [21] G. N. Watson, The final Problem: An account of the mock theta functions, Journal of the London Math, Soc. 11(1936), 55-80.
- [22] G. N. Watson, The Mock Theta Functions (2), Proc. London Math, Soc. 42 (1937), 274-304.
- [23] D. P. Shukla and M. Ahmad, Bilateral mock theta functions of order seven, Math Sci. Res. J. 7(1) (2003), 8-15.
- [24] D. P. Shukla and M. Ahmad, on the behaviour of bilateral mock theta functions of order seven, Math Sci. Res. J. 7(1)(2003), 16-25.
- [25] D. P. Shukla M. Ahmad, Bilateral mock theta functions of order eleven, Proc. Jang Jeon Math Soc. 6(1) (2003), 59-64.
- [26] D. P. Shukla and M. Ahmad, Bilateral mock theta functions of order thirteen, Proc. Jang Jeon Math Soc. 6(2)(2003), 167-183.
- [27] D. P. Shukla and M. Ahmad, Bilateral mock theta functions of order $2r+1$ and their behaviour on the unit circle, "GANITA", Journal of Bharata Ganita Parisad, Vol 60, No. 1, (2009), 27-56.
- [28] Clifford Truesdell. An essay toward a unified theory of special functions, 18 Princeton University Press, (1948).
- [29] Bhaskar Srivastava, Certain bilateral basic hypergeometric transformations and mock theta functions, Hiroshima Maths J. 29 (1999), 19-26.

[30] Bhaskar Srivastava, A study of bilateral forms of the mock theta functions of order eight, Journal of the Chungcheon Math Soc. 18(2)(2005), 117-129.

[31] Bhaskar Srivastava, A mock theta function of second order, International Journal of Mathematics and Mathematical Sciences 2009 (2010), 1-15.

[32] Bhaskar Srivastava, A study of bilateral new mock theta functions, American journal of Mathematics and Statistics 2(4) (2012), 64-69.

[33] B. Gordon and R. J. McIntosh, Some eight order mock theta functions, Journal of the London Math, Soc. 62(2) (2000), 321-335.

[34] Anju Gupta, on certain Ramanujan's mock theta functions, Proc. Indian Acad. Sci. 103(1993), 257-267.