

KINETIC STUDY OF THE SOLVENT EFFECT OF AQUO-TERTIARY ALCOHOL SYSTEMS ON THE CATALYSED SOLVOLYSIS OF BUTYL ACETATE

KAVITA KUMARI,¹R.T. SINGH²

1 [Research Scholar, Dept of Chemistry, V.K.S. University, Ara](#)

2 [Professor & Formerly HOD Chemistry and The Dean of the faculty by Science, V.K.S. university, Ara](#)

ABSTRACT

The solvent effect of a tertiary alcohol was studied on the thermodynamic extensive properties of solvolysis of a heavy acetate (butyl acetate) in different aquo-butanol reaction media containing 20 to 80% (t-butanol) at different temperatures varying from 20 to 40°C. From the enhancement observed in ΔH^ and ΔS^* of the reaction, it is inferred that the organic co-solvent t-butanol acts as entropy controller and enthalpy stimulator solvent for alkali catalysed hydrolysis of butyl acetate. The numerical value of Iso-kinetic temperature of the reaction which comes to be nearly 287.0 (below 300) indicates that there is weak but considerable solvent-solute interaction in the aquo-t-butanol reaction media.*

KeyWords:- *Tertiary butanol, Heavy Acetate, dielectric and Solvation effect. Solvation and desolvation, specific solvation, Extensive properties, Enthalpy Stimulated.*

INTRODUCTION :

Though a large number of kineticists^{1,2} have reported the effect of different solvents on the rates, mechanism and the thermodynamic properties of hydrolysis of simple esters but, a little attention has been paid towards the studies of solvent effect of tertiary alcohol on the rate, mechanism and thermodynamic properties of hydrolysis of simple esters, but, a little attention has been paid towards the studies of solvent effect of tertiary alcohol on the rate, mechanism and thermodynamic activation parameters of the reaction and solvent-solute interaction for alkali catalysed solvolysis of heavy esters. In order to highlight the above

noted facts, it has been proposed to study the kinetics of alkali catalysed hydrolysis of butyl acetate in aquo-t-butanol reaction media.

Experimental & Calculation

The kinetics of alkali catalysed hydrolysis of butyl acetate was carried out separately in the different aquo-organic co-solvent media(aquo-t-butanol) prepared by adding different volumes of t - butanol (20 to 80%). The strength of the solution was kept 0.1M with respect of NaOH and 0.05 M with respect to the ester. The reaction was found to follow the second order kinetic equation and the evaluated values of specific rate constants have been recorded in Table -I. For studying the effect of change of concentration of organic component (t-butanol), The variation of log k values of the reaction with mol% of t-butanol in the reaction media has been enlisted in Table-II. Using Arrhenius equation, the iso-composition and iso-dielectric activation energies values of the reaction were evaluated and are recorded respectively in Table -III & IV. The thermodynamic activation parameters such as ΔH^* , ΔG^* and ΔS^* have been evaluated using Wynne-Jones and Eyring³ equation and their consolidated values have been shown in Table-V. For studying the mechanism of the reaction, the evaluated number of water molecules associated with the transition state of the reaction at different temperatures have been depicted in Table-VI.

Results and Discussion

Effect of solvent on the specific rate constants of the reaction:

In order to highlight the effect of the solvent on the specific rate constant values of the reaction, the log k values were plotted against the mole% of the organic co-solvent (their values from Table-II) as shown in Fig. -1, and were found to follow decreasing trends. However, the depletion found in the rate with increasing mole% of the organic co-solvent (t-butanol) at all the temperatures follow smooth path following two intersection straight lines at about 18.01 mol% of t-butanol having different numerical values of the slope (of similar nature) before and after the point of intersection (at about 18.10mol% of t-butanol in the reaction media). From the plots, it was found that with increasing the temperature of the

reaction, the degree of depletion in the rate constants of the reaction becomes steeper.

However, the possible rate depleting factors in the rate can be listed as follows:

- (i) decrease in the bulk dielectric constant value of the medium,
- (ii) decrease in the polarity of the reaction media on adding less polar t-butanol to it

The above noted two depletion factors are quite in operation and this is quite in agreement with the theory of Hughes and Ingold⁴ that the rate ought to decrease with decreasing dielectric constant of the reaction media. Such decrease in rate constant with increasing proportion of the organic co-solvent like t-butanol has been reported earlier by Singh & Jha et al⁵, Sing & Hafizee et al.⁶ and in recent years also by R.T Singh.⁷ However, the decrease observed in the specific rate constant values with different numerical values of slopes may be attributed partly due to the dielectric effects of the reaction media and partly due to solvation changes taking place in it (aquo-t-butanol reaction media.)

Effect of Solvent on the Iso-composition Activation Energy (E_C) of the Reaction:

On perusal of the data mentioned in Table-III, it is observed that the value of iso-composition activation energy of the reaction go on decreasing from 97.07 kJ/mol to 62.45 kJ/mol with increasing concentration of t-nutanol from 20 to 80% (v/v), in the reaction media. The depletion E_C values of the reaction in water-t-butanol media may be due to either of the following three causes:

- (i) The transition state is more solvated than the initial state,
- (ii) The transition state is les desolvated than the initial state, and
- (iii) The transition state is solvated and the initial state is desolvated.

Among these three factors, the third factor seems to be operative in this case as from the values of thermodynamic activation parameters of the reaction in Table - V, both ΔH^* and ΔS^* values of the reaction are found to decrease with increasing proportion of t-nutanol in the reaction media(ΔH^* values decreases from 92.31 kJ/mol to 60.44 kJ/mol and ΔS^* values

decreases from 30.27 J/K/mol to -81.94 J/K/mol at 30°C. Regarding effect of solvent on the E_C values of reaction, similar findings and their interpretations have been found in accordance with the earlier reports reports of Priyanka & Singh et al⁸. and also with the recently reported findings of Abhay & Singh et al⁹.

Solvent Effect on the Iso-dielectric energy (E_D) of the reaction:

From the values recorded in Table-IV, it appears that E_D values of the reaction go on increasing with increasing dielectric constant values of the aquo-t-butanol reaction media. The E_D values is 78.89 kJ/mol at D values 25 and increases to 98.03 kJ/mol at D value 55. The enhancement in the E_D values with increase in D values of the reaction media is in accordance with depletion in E_C or E_{exp} values of the reaction with increasing mol % of t-butanol in the reaction media. Theses findings and conclusions have been found in support of the earlier reports of Woldford¹⁰ and also with the recent reportings of Tauheed & Singh et al¹¹

Solvent Effect on Thermodynamic Activation Parameters of the Reaction:

From Table -V, on perusal of the values of htermodynamic activation parameters, namely ΔG^* , ΔH^* and ΔS^* , it is observed that ΔG^* values (free energy of activation) of the reaction increases with simultaneous decrease in its ΔH^* and ΔS^* values. At 30°C, ΔG^* values have been observed increasing from 83.14 k cal/mol to 85.27 k cal/mo with increasing concentration of t-butanol from 20 to 80% (v/v) in the reaction media. Though this enhancement is not very large, however, it is quite considerable and acceptable too. In order to highlight the effect of changing concentration of the organic content (t-butanol) in the reaction media, ΔH^* , ΔG^* and ΔS^* values have been plotted against the changing mol% of t-butanol in the reaction media and their plots are shown in Fig. -2, 3 and 4 respectively. From the plots of ΔG^* values against mol % of t-butanol as shown in Fig.-3, it is found that ΔG^* values go on increasing non-linearly with gradual addition of t-butanol in the reaction media. This finding is indicative of desolvation of reactants as explained by Elsemongy et al.¹²

So far as the variations in ΔH^* and ΔS^* are concerned on observing their values from Table-V and their plots against mol % of t-butanol as shown in Fig. -2 and 4, it is

interestingly found that both of them decrease linearly and non-linearly respectively with gradual addition of t-butanol in the reaction media.

From the thermodynamic relation:

$$\Delta G^* = \Delta H^* - T\Delta S^*$$

it can be easily concluded that increase in ΔG^* values with simultaneous decrease in both of ΔH^* and ΔS^* values is only possible when ΔS^* values decreases more than ΔH^* value. From such findings, it is inferred that in presence of t-butanol in the reaction media, the alkali catalysed hydrolysis of butyl acetate becomes entropy controlled and enthalpy stimulated reaction. Moreover, linear variation in ΔH^* and non-linear variation in ΔS^* values with increasing mol% of t-butanol as shown in Fig.-2 and 4 respectively gives information of the fact that specific solvation is taking place in aquo-t-butanol solvent systems similar to that as reported in the past by Saville et al.¹³

However, such solvent effect on thermodynamic activation parameters and extensive properties and their interpretations have also been found in support of the earlier reports of Singh & Priyanka et al.¹⁴ and also with the recent reports of Sushma & Singh et al¹⁵

Solvent Effect on Iso-kinetic Temperature and Solvent-Solute Interaction in the

Reaction Media :

The values of the iso-kinetic temperature of the reaction was evaluated by using Barclely-Butler¹⁶ relationship which is expressed as

$$\delta_m(\Delta H^*) = \beta \delta_m(\Delta S^*)$$

It is a straight line equation representing the relationship between enthalpy and entropy of activation values of the reaction. ' β ' is known as iso-kinetic temperature. From the values of ΔH^* and ΔS^* values available in Table -V. the plots of ΔH^* versus ΔS^* at 30°C were made which is shown in Fig. -5. From the slope of the straight line of the plots, the values of the kinetic temperature was evaluated to be $287.32 \approx 287.0$ (below 300). Thus, in the light of Leffler's guidelines¹⁷, from the numerical values of the iso-kinetic temperature

(which is below 300), it can safely be concluded that there is appreciable change in the structure of the reactant or in the solvent or in both the reactant and solvent due to weak but considerable interaction between solvent and solute present in the reaction media (aquo-t-butanol) in the similar way as reported earlier by Singh & Singh et al.¹⁸ and also in recent years by Amant & Singh et al.¹⁹

Effect of Solvent (water concentration) on the Mechanism of the Reaction:


Evaluation of the number of water molecules associated with the formation of transition state or activated complex has been done by plotting log k values of the reaction against log[H₂O] values as shown in Fig. -6 in the light of relation proposed by Robertson²⁰.

$$\log k = \log k_0 + n \log [\text{H}_2\text{O}]$$

From Fig. -6, it is apparent that at each temperature two straight lines intersecting each other at log[H₂O] value approximately 1.453 having different values of slopes are obtained. From the slopes of the plots of log k versus log [H₂O] the evaluated values of number of water molecules associated with the activated complex of the reaction are noted in Table-VI. From Fig. -6 as well as from Table-VI, it is clear that the number water molecules associated with the transition state increases from 0.236 to 0.771 with increase in temperature above 20 to 40°C before log [H₂O], values 1.453 which corresponds to 51.30% of water concentration in the reaction media.

Similarly, in case of above 51.30% of water concentration in the reaction media, the number of water molecules associated with the transition state in its formation increases from 0.520 to 1.145 with rise of temperature from 20 to 40°C. Overall, the number of water molecules involved in the formation of the activated complex of the reaction increases from 0.236 to 1.145 with rise of temperature from 20 to 40°C. According to observation and findings of Robertson et al.²¹ it has been suggested that the number of water molecules associated with the transition state is fairly high for unimolecular reaction while that for bimolecular reaction will be very low. Hence in the light of findings of Robertson et al.²¹, it may be inferred that with rise in temperature of the reaction, the mechanistic path of the reaction changes from bimolecular to unimolecular in aquo-t-butanol media. From above noted

findings about the increasing number of water molecules associated with activated complex, it may be inferred that in presence of t-butanol in the reaction media and with rise of temperature of the reaction, the structure of water is changes from its bulky form to dense form at equilibrium.



Earlier Akanksha & Singh et al.²² and recently Sushma-Abhay & Singh et al.²³ have also reported similar findings and their interpretations for effect of solvent on the mechanism of the catalysed solvolysis reactions.

References:

1. Elsemongy, M.M. Abu Elamayam, M.S. and Moussa, M.N.H. : Z. Physik Che. (Neue Folge), 95,215, 1975
2. Singh, Lallan, Gupta, A.K., Singh, R.T., Verma, D.K. and Jha, R.C. : React. Kinet. Catal Lett. 24, No(1-2), 161-165, 1984
3. Wynne Jones W.F.K. and Eyring, H. : J. Chem. Phys., 3, 492, 1935
4. Hughes E.D. and Ingold C.K. : J. Chem. Soc. 244, 255, 1935
5. Singh, Lallan, Singh, R.T., and Jha, R.C. : J. Indian Chem. Soc. **58**, 966, , 1981
6. Singh, R.T., Navendu, K.S., Henry, W. and Hafizee, N.K. : NIRJ Sci. 14, 53-61, 2014.
7. Singh, R.T. : ARJ Phys. Sci., **18**, No. (1-2), 105-116, 2015
8. Priyanka, K., Nazia, S., Kumar, V. and Singh, R.T. : ARJ Phys. Sci., 17, No.(1-2), 117-128, 2014
9. Abhay A, Singh R and Singh R.T : NIRJ Sci **22**, 53-66, 2016(Dec)
10. Wolford, R.K. : J. Phys. Chem., **64**, 3392, 1964

11. Tauheed S. Haider R and Singh RT : NIRJ Sci 24, 85-100, 2017(Sept)
12. Elsemongy, M.M Abu Elamayem, M.S. and Moussa, M.N.H. : Z. Physik. Chem. Neue Folge, 84, 295, 1975
13. Saville, B.J. and Husdan, R.F. : J. Chem. Soc., 4114, 1955
14. Singh, R.T., Priyanka, K., Singh, S. and Singh, R.K. :NIRJ Sci. **16**, 15-24, 2014
15. K Sushma Ranjan R K and Singh R T : IJSEM, 7, No(2), 147-162, 2018(Feb)
16. Barclay, I.A. and Butler, J.A.V. : Trans Faraday Soc., 34, 1445, 1938
17. Leffler, J.E. : J. Org. Chem., **20**, 1201, 1955
18. Singh, R.T., Singh, P.K., Singh, S.M. and Singh, U.C. : ARJ Phys. Sci., **15**, No. (1-2), 129-139, 2012
19. Amant and Singh RT : NIRJ Sci **27**, 1-12, 2018(June)
20. Robertson, R.E. : Prog. Phy. Org., Chem. **4**, 213, 1967
21. Robertson, R.E., Hippolittle, R.L. and Scott, J.M.W. : Canad. J. Chem. Soc., **37**, 303, 1959
22. Akanksha, Kuamri, R., Kumar, R. and Singh, R.T. : ARJ Phys. Sci., **17**, No.(1-2), 105-116, 2014
23. K. Sushma Abhay A, Singh R and Singh RT : JICS, 10 No(5), 794-813, 2020

Table - I

**Specific rate constant values of Alkali catalysed hydrolysis of Butyl acetate
in water-t-butanol media
 $K \times 10^2$ in $(\text{dm})^3 \text{mole}^{-1} \text{min}^{-1}$**

| Temp in °C | % of t-butanol (v/v) | | | | | | |
|------------|----------------------|--------|--------|--------|--------|--------|--------|
| | 20% | 30% | 40% | 50% | 60% | 70% | 80% |
| 20 °C | 49.23 | 45.85 | 42.54 | 40.37 | 38.04 | 36.33 | 33.10 |
| 25 °C | 93.80 | 86.18 | 79.03 | 70.42 | 64.24 | 58.05 | 50.13 |
| 30 °C | 177.01 | 158.64 | 144.18 | 123.94 | 108.24 | 95.85 | 75.88 |
| 35 °C | 323.30 | 285.30 | 252.81 | 208.88 | 175.95 | 142.33 | 112.18 |
| 40 °C | 602.70 | 504.89 | 442.69 | 354.65 | 286.26 | 208.22 | 166.05 |

Table - II

Variation of log k values of the reaction at different temperatures with mol % of t-butanol in water-t-butanol media.

| % of t-butanol(v/v) | Mol % of t-butanol | 2+ log k values | | | | |
|---------------------|--------------------|-----------------|--------|--------|--------|--------|
| | | 20°C | 25°C | 30°C | 35°C | 40°C |
| 20% | 4.53 | 1.6922 | 1.9722 | 2.2480 | 2.5096 | 2.7801 |
| 25% | 7.52 | 1.6613 | 1.9354 | 2.2004 | 2.4553 | 2.7032 |
| 30% | 11.23 | 1.6288 | 1.8978 | 2.1589 | 2.4028 | 2.6461 |
| 40% | 15.95 | 1.6061 | 1.8478 | 2.0932 | 2.3199 | 2.5498 |
| 50% | 22.15 | 1.5802 | 1.8078 | 2.0344 | 2.2454 | 2.4568 |
| 60% | 30.68 | 1.5603 | 1.7638 | 1.9678 | 2.1533 | 2.3185 |
| 70% | 43.15 | 1.5198 | 1.7001 | 1.8801 | 2.0499 | 2.2202 |

Table - III

Evaluated values of Iso-composition Activation Energy (E_C or E_{exp}) of the reaction in water-t-butanol media.

| % of t-butanol(v/v) | 20% | 30% | 40% | 50% | 60% | 70% | 80% |
|-------------------------|-------|-------|-------|-------|-------|-------|-------|
| E_C values in K/J/mol | 97.07 | 91.41 | 89.50 | 83.30 | 77.39 | 67.86 | 62.45 |

Table - IV

Evaluated Values of Iso-Dielectric Activation Energy (E_D) of the reaction at Desired 'D' values of water-t-butanol media.

| D values | D= 32 | D = 37 | D = 42 | D =47 | D = 52 | D = 57 | D = 62 |
|-------------------------|-------|--------|--------|-------|--------|--------|--------|
| E_D values in kJ/mole | 78.89 | 82.38 | 84.83 | 88.25 | 90.01 | 92.81 | 95.03 |

Table - V

Values of the slopes of the plots of $\log k$ versus $\log [H_2O]$ values at different temperatures

| Temperature in °C | Slope - I | Slope - II |
|-------------------|------------------------------------------|-----------------------------------------|
| | Where $\log [H_2O]$ value is below 1.453 | when $\log [H_2O]$ value is above 1.453 |
| 20°C | 0.236 | 0.520 |
| 25°C | 0.434 | 0.692 |
| 30°C | 0.552 | 0.794 |
| 35°C | 0.725 | 0.956 |
| 40°C | 0.771 | 1.145 |

Table- VI

Consolidated Values of Thermodynamic Activation Parameters (ΔH^* , ΔG^* and ΔS^*) of the reaction in water-butanol system at different temperatures.

ΔH^* and ΔG^* in kJ/mol, ΔS^* in J/K/mol

| % of t-butanol (v/v) | Mole% of t-butanol | ΔH^* in kJ/mol | 20°C | | 25°C | | 30°C | | 35°C | | 40°C | |
|----------------------|--------------------|------------------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | | | ΔG^* | ΔS^* | ΔG^* | ΔS^* | ΔG^* | ΔS^* | ΔG^* | ΔS^* | ΔG^* | ΔS^* |
| 20% | 4.53 | 92.31 | 83.43 | 30.30 | 83.29 | 30.24 | 83.14 | 30.27 | 83.02 | 30.14 | 82.78 | 30.46 |
| 30% | 7.52 | 89.55 | 83.60 | 20.29 | 83.51 | 20.27 | 83.41 | 20.24 | 83.33 | 20.19 | 83.24 | 20.16 |
| 40% | 11.23 | 87.19 | 83.78 | 11.62 | 83.72 | 11.43 | 83.65 | 11.67 | 83.64 | 11.53 | 83.58 | 11.54 |
| 50% | 15.95 | 80.37 | 83.91 | -12.07 | 84.01 | -12.18 | 84.03 | -12.08 | 84.13 | -12.18 | 84.16 | -12.08 |
| 60% | 22.15 | 74.69 | 84.06 | -31.97 | 84.23 | -32.03 | 84.37 | -31.96 | 84.57 | -32.06 | 84.71 | -32.02 |
| 70% | 30.68 | 65.45 | 84.17 | -63.90 | 84.49 | -62.83 | 84.76 | -63.75 | 84.11 | -63.84 | 85.54 | -64.20 |
| 80% | 43.15 | 60.44 | 84.39 | -81.75 | 84.85 | -80.38 | 85.27 | -81.94 | 85.72 | -82.07 | 86.13 | -82.00 |

